

Introduced in Great Britain in 1949 for ornamental purposes. The birds escaped from captivity and reproduced for the first time in 1960. They arrived in France in 1974.

20	cri	nı	H
CO	UI I	P	
		•	

- A small, diving duck 35 to 45 cm long, with a wingspan of 53 to 62 cm
- The ducks have a large tail, short wings and a large bill
- Clear sexual dimorphism:
- the average weight of females is 510 grams, of males 610 grams
- breeding males are ruddy brown in colour with white cheeks and black top of head, a sky blue bill and white under the tail. Their eclipse plumage is greyish brown, but the head is identical
- females and immature males are a drab brown with a brown bill and yellowish cheeks with a brown stripe


## **Ecology and reproduction**

- Habitats include:
- water bodies with dense riparian vegetation
- fresh or brackish water, however in France, they are found exclusively in fresh waters
- The animals may be aggressive with congeners and other species during the breeding season, but are peaceable otherwise
- Their diet consists of aquatic plants, molluscs, crustaceans, worms and insects
- Reproduction occurs from June to the end of August:
- during the period, the animals are highly dispersed with their nests hidden in the riparian vegetation
- broods count 6 to 14 eggs that incubate for 25 days
- the young take off on their own after 50 to 55 days
- first reproduction at age of 2 years

## **D**ocumentation

- Sarat E. (coord.) 2012. Vertébrés exotiques envahissants du bassin de la Loire (hors poissons) : connaissances et expériences de gestion. Office national de la chasse et de la faune sauvage, Plan Loire Grandeur Nature, 128 pp.
- Madge S., Burn H. 1995. Guide des canards, des oies et des cygnes. Delachaux et Niestlé, Lausanne. 304 pp.
- Caizergues A., Fouque C. 2008. Zoom sur l'Érismature rousse, une espèce à éradiquer en France. Faune Sauvage, 280 : 64-66.

Author: Emmanuelle Sarat, IUCN French committee









- 1. Male ruddy duck with breeding plumage.
- 2. Female ruddy duck.
- 3. Aggressive males during the breeding season.

